

E11 Upper GI Endoscopy and Colonoscopy

Expires end of March 2024

If you have any questions or concerns about your care, please contact the nurse in charge at the hospital.

You can get more information locally from:

You can also get information from www.aboutmyhealth.org
Tell us how useful you found this document at www.patientfeedback.org



Royal College
of Surgeons
of England



THE ROYAL
COLLEGE OF
SURGEONS
OF EDINBURGH



Patient Information Forum

eidohealthcare.com



UNITED KINGDOM

What is an upper GI endoscopy and colonoscopy?

An upper gastrointestinal (GI) endoscopy is a procedure to look at the inside of your oesophagus (gullet), stomach and duodenum using a flexible telescope. This procedure is sometimes known as a gastroscopy, OGD or simply an endoscopy.

A colonoscopy is a procedure to look at the inside of your large bowel (colon) using a flexible telescope.

Your doctor has suggested an upper GI endoscopy and colonoscopy. However, it is your decision to go ahead with the procedure or not. This document will give you information about the benefits and risks to help you to make an informed decision.


If you have any questions that this document does not answer, it is important that you ask your doctor or the healthcare team. Once all your questions have been answered and you feel ready to go ahead with the procedure, you will be asked to sign the informed consent form. This is the final step in the decision-making process. However, you can still change your mind at any point.

What are the benefits of an upper GI endoscopy and colonoscopy?

Your doctor is concerned that you may have a problem in your digestive system. They may have recommended an upper GI endoscopy and colonoscopy because you have been experiencing symptoms such as stomach pain, difficulty swallowing, bloating, diarrhoea and/or unexplained weight loss. Another reason might be that you do not have symptoms, but you are anaemic. This procedure is a good way of diagnosing most problems with your digestive system.

If the endoscopist (the person doing the endoscopy and colonoscopy) finds a problem, they can perform biopsies (removing small pieces of tissue) to help make the diagnosis.

Sometimes a polyp (small growth) is the cause of the problem and the endoscopist may be able to remove it during the procedure.



An upper GI endoscopy

Are there any alternatives to an upper GI endoscopy and colonoscopy?


Your doctor has recommended an upper GI endoscopy and colonoscopy as it is the best way of diagnosing most problems with your digestive system.

A barium meal is an x-ray test of your upper digestive system. This test is not as accurate as an upper GI endoscopy. Alternatives to a colonoscopy include a CT colography (a CT scan of your large bowel).

If your doctor finds a problem, you may still need an upper GI endoscopy or colonoscopy to treat the problem or perform biopsies.

What will happen if I decide not to have an upper GI endoscopy and colonoscopy?

Your doctor may not be able to confirm what the problem is. If you decide not to have an upper GI endoscopy and colonoscopy, you should discuss this carefully with your doctor.



A colonoscopy

What does the procedure involve?

Before the procedure

If you take iron tablets, stop taking them at least a week before the procedure.

If you take warfarin, clopidogrel or other blood-thinning medication, let the endoscopist know at least 7 days before the procedure.

The healthcare team will give you instructions about when you need to stop eating and drinking to make sure the endoscopist can have a clear view of your stomach and colon.

You will need to follow a special diet and you will be given some laxatives to take the day before the procedure. Follow the instructions carefully.

If you have diabetes, let the healthcare team know as soon as possible. You will need special advice depending on the treatment you receive for your diabetes.

If you get severe abdominal pain or if you vomit, contact the endoscopy unit or your doctor.

The procedure may involve injecting you with medication (Buscopan) to relax your bowel and make the procedure more comfortable. Buscopan can affect the pressure in your eyes, so let the endoscopist know if you have glaucoma.

The healthcare team will carry out a number of checks to make sure you have the procedure you came in for. You can help by confirming to the endoscopist and the healthcare team your name and the procedure you are having.

The healthcare team will ask you to sign the consent form once you have read this document and they have answered your questions.

In the endoscopy room

An upper GI endoscopy and colonoscopy usually takes about an hour.

The endoscopist may offer you a sedative or painkiller to help you to relax. They will give it to you through a small needle in your arm or the back of your hand. You will be able to ask and answer questions but you will feel relaxed. You may not be aware of or remember the procedure. The healthcare team can give you more information about this.

Many endoscopists perform the procedure with you asleep under a general anaesthetic, which they can give you through the small needle, or as a mixture of anaesthetic gas that you breathe through a mask or mouthpiece. This means you will be unaware of the procedure. The healthcare team can give you more information about this.

Once you have removed any false teeth or plates, they may spray your throat with some local anaesthetic and ask you to swallow it. This can taste unpleasant.

The endoscopist will ask you to lie on your left side and will place a plastic mouthpiece in your mouth. The healthcare team will monitor your oxygen levels and heart rate using a finger or toe clip. If you need oxygen, they will give it to you through a mask or small tube under your nostrils.

If you are awake during the procedure and at any time you want it to stop, let the endoscopist know. The endoscopist will end the procedure as soon as it is safe to do so.

An upper GI endoscopy involves placing a flexible telescope (endoscope) into the back of your throat. The endoscopist may ask you to swallow when the endoscope is in your throat. This will help the endoscope to pass easily into your oesophagus and down into your stomach. From

here the endoscope will pass into your duodenum.

A colonoscopy involves placing a flexible telescope into your back passage and blowing some air into your large bowel to get a clear view.

The endoscopist will be able to look for problems such as inflammation, ulcers or polyps (small growths). They will be able to perform biopsies and take photographs to help make the diagnosis. If they find a polyp, it may be possible to remove it during the procedure.

What complications can happen?

The healthcare team will try to reduce the risk of complications.

Any numbers which relate to risk are from studies of people who have had this procedure. Your doctor may be able to tell you if the risk of a complication is higher or lower for you.

Some complications can be serious and can even cause death (risk: less than 3 in 25,000).

You should ask your doctor if there is anything you do not understand.

The possible complications of an upper GI endoscopy and colonoscopy are listed below.

- Sore throat. This gets better quickly.
- Breathing difficulties or heart irregularities, as a result of reacting to the sedative, inhaling secretions such as saliva, or your bowel being stretched. To help prevent this, your oxygen levels and heart rate will be monitored and a suction device will be used to clear any secretions from your mouth.
- Heart attack (where part of the heart muscle dies) or stroke (loss of brain function resulting from an interruption of the blood supply to your brain) can happen if you have serious medical problems. This is rare.
- Allergic reaction to the equipment, materials or medication. The healthcare team is trained to detect and treat any reactions that might happen. Let the endoscopist know if you have any allergies or if you have reacted to any medication or tests in the past.
- Infection. It is possible to get an infection from the equipment used, or if bacteria enter

your blood. The equipment is disinfected so the risk is low but let the endoscopist know if you have a heart abnormality or a weak immune system. You may need treatment with antibiotics. Let your doctor know if you get a high temperature or feel unwell.

- Blurred vision, if you are given a Buscopan injection. This usually gets better after about an hour. Sometimes the injection can also affect the pressure inside your eye. This is more likely if you have a rare type of glaucoma. If your eye becomes red and painful, and your vision becomes blurred, let your doctor know straight away.
- Making a hole in your oesophagus, stomach, duodenum or colon (risk: less than 3 in 2,000). The risk is higher if a polyp is removed, especially if it is a large polyp. This is a serious complication. You may need surgery which can involve forming a stoma (your bowel opening onto your skin).
- Damage to teeth or bridgework. The endoscopist will place a plastic mouthpiece in your mouth to help protect your teeth. Let the endoscopist know if you have any loose teeth.
- Bleeding from a biopsy site or from minor damage caused by the endoscope (risk: less than 1 in 1,000). This usually stops on its own.
- Bleeding, if a polyp is removed (risk: 2 in 100). This may be higher if you have multiple or large polyps removed. Bleeding usually stops soon after a polyp is removed. Sometimes bleeding can happen up to 2 weeks after the procedure. If you take blood-thinning medication and have a polyp, the endoscopist will usually not remove it.
- Missed polyp. Let your doctor know if you have any problems with your bowel after the colonoscopy.
- Incomplete procedure caused by a technical difficulty, food or blockage in your digestive system, complications during the procedure, or discomfort. Your doctor may recommend another endoscopy or colonoscopy, or a different test such as a barium meal or CT colography.

How soon will I recover?

After the procedure you will be transferred to the recovery area where you can rest. If you were given a sedative, you will usually recover in about 2 hours but this depends on how much sedative you were given.

Do not eat or drink for at least the first hour after the procedure. Once you are able to swallow properly, you will be given a drink. You may feel a bit bloated for a few hours but this will pass.

If you had sedation:

- a responsible adult should take you home in a car or taxi and stay with you for at least 24 hours;
- you should be near a telephone in case of an emergency;
- do not drive, operate machinery or do any potentially dangerous activities (this includes cooking) for at least 24 hours and not until you have fully recovered feeling, movement and co-ordination; and
- do not sign legal documents or drink alcohol for at least 24 hours.

You should be able to return to work the next day unless you are told otherwise.

The healthcare team will tell you what was found during the procedure and discuss with you any treatment or follow-up you need. Results from biopsies will not be available for a few days so the healthcare team may arrange for you to come back to the clinic for these results.

Once at home, if you get pain in your chest or abdomen, difficulty breathing, significant or continued bleeding from your back passage, a high temperature, or if you vomit, contact the endoscopy unit or your GP. In an emergency, call an ambulance or go immediately to your nearest Emergency department.

Lifestyle changes

If you smoke, stopping smoking will improve your long-term health.

Try to maintain a healthy weight. You have a higher risk of developing complications if you are overweight.

Regular exercise should improve your long-term health. Before you start exercising, ask the healthcare team or your GP for advice.

Summary

An upper GI endoscopy and colonoscopy is usually a safe and effective way of finding out if there is a problem with your digestive system. However, complications can happen. You need to know about them to help you to make an informed decision about the procedure. Knowing about them will also help to detect and treat any problems early.

Keep this information document. Use it to help you if you need to talk to the healthcare team.

Some information, such as risk and complication statistics, is taken from global studies and/or databases. Please ask your surgeon or doctor for more information about the risks that are specific to you, and they may be able to tell you about any other suitable treatments options.

This document is intended for information purposes only and should not replace advice that your relevant healthcare team would give you.

Acknowledgements

Reviewers

Simon Parsons (DM, FRCS)

Jonathan Lund (DM, FRCS)

Illustrator

Medical Illustration Copyright ©
Medical-Artist.com